

A CONTRIBUTION TO THE FLORA OF TUWA IN PANCHMHAL DISTRICT GUJARAT¹

G. L. SHAH AND M. B. DESHPANDE

Saurashtra Poly. University, Rajkot (Gujarat)

ABSTRACT

The areas in vicinity of Tuwa village have been intensively explored from 1964-1968. During this period 329 angiosperms were collected of which 209 are dicotyledons and 120 monocotyledons. There are five new records for vertebral Bombay Presidency and one new species. The vegetation is dominantly a semi-evergreen type but some components of dry deciduous forests are also met with. A comparison is made with that of Pavagadh hill with a view to bring out differences between the two areas which are topographically different.

INTRODUCTION

The flora of some forests in the hilly regions of Panchmahal district such as Lonavada (Saxxon, 1922), Desighdharia (Saxxon) and Ramanandpur (Chavan and Bedi, 1966; Bedi, 1968) is known to some extent but it is better known for Pavagadh hill (Shah and Inamdar 1953; Chavan and Oza, 1966; Bedi, Sabnis and Bharat 1968; Chokan and Shah, in press). However, it seems that no such work is undertaken in the plains of the district. A study of this type would be helpful in comparing the flora of the hilly regions with that of plains. With this end in view the authors studied the flora of Tuwa and its environs. The selection of the area was made as a matter of convenience. Intensive explorations in different seasons of the year were carried out from 1964 to 1968. We have been able to record 5 new plants for the erstwhile Bombay Presidency and 1 new species (Chokan, Shah and Deshpande, 1967; Shah and Deshpande, 1968; Deshpande and Shah, 1968). About 339 species have been listed in this paper of which 209 are dicotyledons and 130 monocotyledons. The nomenclature adopted is the one published in the current authentic taxonomical literature. The herbarium specimens, which serve as a basis for this paper, are housed in the Botany Department, S. P. University, Vallabh Vidyanagar.

GEOGRAPHY AND CLIMATE OF THE AREA

Tuwa is a small village in N. W. part of Panchmahal district in eastern Gujarat, situated $22^{\circ}45'N$ and $73^{\circ}27'E$ at an elevation of about 56 m above Mean Sea Level. It is a small railway station on

Anand-Godhra line of the Western Railway about 30 km south of Godhra. It is also approachable by the State Transport buses which regularly ply between Anand and Godhra along the highway which runs within the limits of the area explored, almost parallel to the railway line. Tuwa is also famous in Gujarat for its hot water springs, which are at a distance of only about one kilometer from Tuwa station. Topographically the area is a plain country traversed by three main rivers—Klat, Moro and Bochad and their tributaries. The soil is mostly alluvial, varying in colour from pale to dark brown. The hottest months are March-May, October being a close second. The rainy season starts in the middle of June or beginning of July and lasts till September. The average annual rainfall is about 697.14 mm. The cold season is from November to February but December-January are the coldest months.

SOME ASPECTS OF THE VEGETATION OF THE AREA

The vegetation of the hilly regions of Panchmahal district is of a dry deciduous type (Saxxon and Sedgwick, p. 228), but in the present area, it is an open and scrub forest. Even there, some trees like *Butea monosperma* Taub. and *Tectona grandis* L.f. which often form a dominant vegetation in dry deciduous forests, are common. The other members which are also frequent components of such forests are but rarely met with. They are *Bombax ceiba* L., *Buchanania laurina* Spr., *Bridelia squamula* Gurm., *Diospyros melanoxylon* Roxb.

¹Based on the Ph. D. Thesis approved by S. P. University, Vallabh Vidyanagar.
Unpublished collection housed in Gujarat College Herbarium, Ahmedabad.

Doliandra latifolia var. *brevia* Blume, *Grewia arborea* Roth, *Ixora brachiata* Roth, *Millettia tomentosa* Sind., *Mitragyna parvifolia* Korth., *Sympodium lehingua* A. Juss., *Sterculia urens* Roth; and *Ternstroemia crenulata* Roth. This may therefore suggest that the vegetation of the present area must have been in the past similar to the one found in the hilly region. However the dominance of thorny trees and shrubs can be attributed to biotic causes operating for the last many years. The local people indiscriminately cut down valuable trees and shrubs for fuel and other domestic purposes. The tree-like *Butes manisperma* Taub., *Tecoma grandis* L. f. etc. are seldom allowed to grow by themselves in many places and, as a result, they remain stunted, only up to 1.5 m tall. Along railway lines vast stretches of land are covered with mixed scattered patches of the stunted forms of these trees. At times *Diospyros melanoxylon* Roth., *Holarhena annulipes* Wall. ex DC., *Maytenus emarginata* Hor., are also found in these patches. Further, the cattle graze at random, causing a number of tree and shrub seedlings and herbs to disappear.

A comparison of the vegetation of the present area with that of the Paragadh hill will not be out of place. Though the vegetation of the hill is of a dry deciduous type, some members of the scrub forest such as *Acacia salicina* esp. *indica* Benth., *Balanites aegyptiaca* Del., *Buchanania coriaria* W. & A., *Maytenus emarginata* Hor., *Ziziphus mucronata* W. & A. ex., which are common in our area, are also found at the foot and on hilly slopes. The dry deciduous nature of the forests of the hill has been preserved because of very much less, or practically no, biotic interference. Chavan and Oza (p. 5) have rightly remarked "The area of this isolated hill has been declared 'unworkable areas'-forest reserve' by the forest department." This is left to the nature and hence the luxuriant growth of the vegetation which as a consequence is not disturbed by fire, ploughing, grazing by sheep and other animals.

Though the area selected by us is rather limited, about 6 km radius from Tiwa railway station, covering some neighbouring villages like Vagiriyar, Kankapur, Vavadihord, Ranaper, Vanjai, Chachpur, Gali and Dhonol, the flora of our area is rich, as can be judged from the following table showing 12 most abundant families in the present area and their relative positions in the flora of

Paragadh. The numbers in parenthesis stand for the number of species represented in a family.

Sl.No.	Name of the family	Flora of the area including deciduous and semi-evergreen areas up to sea level 200 m	Paragadh, hill elevation 720 m above the sea level, including plains
1.	Lamiaceae	173	160
2.	Cannabaceae	273	270
3.	Cyperaceae	3642	238
4.	Compositae	5052	230
5.	Convolvulaceae	2023	223
6.	Malvaceae	613	213
7.	Euphorbiaceae	2119	227
8.	Scrophulariaceae	819	114
9.	Acanthaceae	219	123
10.	Poaceae	1012	306
11.	Oleaceae	1111	815
12.	Labiaceae	1219	514

Total number of plants in the present area — 529
 Total number of plants in Paragadh —
 So far reported or known — 590
 Cultivated plants in the present area — 20 approximately
 Cultivated plants in Paragadh flora — 20 approximately 3.5%

From the above table it will be seen that the first eleven most abundant families represented in Paragadh flora are also comparatively better represented in our area, but the differences in the number of species in each family are obviously due to topography of the two areas and the biotic influence in the present area. For example the less number of grasses in Tiwa and its environs may be due to indiscriminate grazing by cattle. On the other hand the Cyperaceae and the other hydrophytes are very well represented in Tiwa but poorly so at Paragadh. This is because the plain country of Tiwa and its environs is traversed by rivers, temporary ponds, padillas, ditches which support a good number of cyperaceous plants along with other hydrophytes. Some of the noteworthy hydrophytes are submerged—*Aponogeton natans* Engl., *Ceratophyllum demersum* L., *Hydrilla verticillata* Rott., *Najas minor* All., *N. guadalupensis* Rendle, *Potamogeton pectinatus* L. and *Kallstroemia speciosa* L.; floating—*Ipomoea aquatica* Forsk., *Lemnophyton antistellatum* Miq., *Ludwigia adscendens* Hara, *Neptria obtusa* Lour., Nymphaeacean members, *Nymphoides cristata* O.K., *N. indica* O.K. etc. A number of amphibious plants including Cyperaceae were noted. A comparison of the flora of Tiwa with that of Paragadh hill will also show that most of the plants are common in both the areas but some of them occurring in our

etc. are not reported from Pavagadhi and vice versa.

ENUMERATION

ANNONACAE

Annona squamosa Linn.
Millettia tomentosa (Roxb.) Sweet

MENISPERMACAE

Cissampelos pareira Linn.
Coccinia hispida (Linn.) Didié
Tinospora cordifolia (Willd.) Miers ex Hook. E. & Thoms.

NYMPHORACEAE

Nelumbo nucifera Gaertn.
Sympetrum nouchali Burm. f.
N. stellata Willd.

PARVIVERBACEAE

Argemone mexicana Linn.

BRASSICACAE CRUCIFERAE

Brassica juncea Coss.
B. nigra (Linn.) Koch.
Lepidium sativum Linn.
Raphanus sativus Linn.

CAPPARACAE

Cathaba fruticosa (Linn.) Druec
Capparis decidua (Forsk.) Edgew.
C. sepiaria Linn.
Oxome beachycarpa Vahl ex DC
C. gynandra Linn.
C. simplicifolia (Cavill) Hook. f. & Thoms.
C. viscosa Linn.
Marcus oblongifolia A. Rich.

VIOLACEAE

Byblanthus enneaspernum (Linn.) F. Muell.

FLACOURTIACAE

Eurycoma longifolia (Burm. f.) Merr.

POLYGALACAE

Polygalia chinensis Linn.
P. triptera DC.

CARYOPHYLLACAE

Polyarpaea corymbosa (Linn.) Lamk.
Polyarpaea prostratum Aschers. & Schweinf.
Spergula arvensis Linn.
Vaccaria pyramidata Medic.

PORTULACACEAE

Petalostoma oleracea Linn.
P. quadrifida Linn.

TAMARICACEAE

Tamarix ericoides Rott.

TILIACEAE

Bergia ammanoides Roxb.
B. odorata Edgew.
Elatine ambigua W.L.

MALVACAE

Abelmoschus esculentus (Linn.) Vierch.
A. ficinicus (Linn.) Wt. & Arn.
A. manihot (Linn.) Medic.
Abutilon indicum (Linn.) Sw.
Gossypium herbaceum Linn.
Hibiscus cannabinus Linn.
H. lobatus (Mars.) O. Kuntze
H. ovalifolius (Forsk.) Vahl
H. sabdariffa Linn.
H. vitifolius Linn.
Pavonia zeylanica (Linn.) Cav.
Sida cordifolia Linn.
S. spinosa Linn.
S. veronicæfolia Lamk.
Thespesia populnea (Linn.) Soland. ex Gou

Urena lobata Linn.

BOMBACACEAE

Bombax ceiba Linn.

STERCULIACEAE

Helicteres isora Linn.
Melochia corchorifolia Linn.
Sterculia urens Roxb.
Waltheria indica Linn.

TILIACEAE

Cochruss acuminatus Linn.
C. capsularis Linn.
C. fascicularis Lamk.
C. olitorius Linn.
Grewia flavescens Juss.
Triumfetta pentandra A. Rich.
T. rhomboidea Jacq.
T. rotundifolia Lamk.

LINACEAE

Linum usitatissimum Linn.

ZYGOPHYLLACEAE

Tribulus terrestris Linn.

OXALIDACEAE

Biophytum sensitivum (Linn.) DC.

RUTACEAE

Aegle marmelos (Linn.) Coll.
Citrus limon (Linn.) Burm. f.

- Feronia limonia* (Linn.) Swingle
Murraya koenigii (Linn.) Sprague
 SIMARUBACEAE
Allanthe excelsa Roxb.
 RALANTACEAE
Batamites aegypticus (Linn.) Del.
 MELIACEAE
Azadirachta indica A. Juss.
Sympidium lebrili A. Just.
 CELASTRACEAE
Celastrus paniculatus Willd.
Maytenus emarginata (Willd.) Dc. Hott
 RILIMINACAE
Zizyphus mauritiana Lamk.
Z. nummularia (Burm. f.) Wt. & Arn.
Z. otocarpa (Linn.) Mill.
Z. xylopyrus (Rech.) Willd.
 VITACEAE
Ampelocissus latifolia (Roxb.) Blanch.
Cayratia carnosia (Lamk.) Gagnep.
 SAPINDACEAE
Cardiospermum halicacabum Linn.
 ANACARDIACEAE
Buchanania lanzen Sprague
Lannea coromandelica (Horn.) Merr.
Mangifera indica Linn.
 MORINGACEAE
Moringa concreta Nairne
M. oleifera Lamk.
 PAPILIONACEAE
Abrus precatorius Linn.
Aesculus indica Linn.
Alysicarpus hirsutus (Linn.) DC.
A. longifolius (Roxb. ex Sprague) Wt. & Arn.
A. monilifer (Linn.) DC.
A. procumbens (Roxb.) Schindl.
A. scorpioides Griseb. ex Thw.
A. tectorum Edgew.
A. vaginalis (Linn.) DC.
Arachis hypogaea Linn.
Atylosia platycarpa Benth.
A. scrophularia (Linn.) Benth.
Butea monosperma (Linn.) Taub.
Cajanus cajan (Linn.) Millsp.
Canavalia gladiata (Jacq.) DC.
Cicer arietinum Linn.
Citorea ternata Linn.
Crotalaria albida Heyne ex Roth
C. burhia Benth. Hau. ex Benth.
C. juncea Linn.
C. linifolia Linn. L.
C. medicaginea Lamk.
C. retusa Linn.
Cynometra tetragonoloba (Linn.) Tand.
Diospyros dichotomum (Willd.) DC.
D. gangeticum (Linn.) Willd.
D. triflorum (Linn.) DC.
Dolichos lablab Linn.
Elettaria monophylla (Burm. f.) DC.
Heylandia latibrosa DC.
Indigofera astragalina DC.
I. cordifolia Heyne ex Roth
I. glandulosa Willd.
I. limitata Retz.
I. linnaei All.
I. oblongifolia Forsk.
I. risterioria Linn.
I. trita Linn. L.
Medicago sativa Linn.
Mellilotus indica All.
Mucuna pruriens Hook. f.
Phaseolus aconitifolius Jacq.
P. angularis (Willd.) Wt.
P. radiatus Linn.
P. trilobus All.
Pongamia pinnata (Linn.) Pierre
Pueraria tuberosa (Roxb. ex Willd.) DC.
Rhynchosia minima (Linn.) DC.
R. sutea DC.
Sesbania hispida (Jacq.) W. F. Wight
Tephrosia purpurea (Linn.) Pers.
T. villosa (Linn.) Pers.
Teramnus labialis (Linn. f.) Spreng.
Trigonella foenum-graecum Linn.
Vigna unguiculata (Linn.) Walp.
Zornia gibbosa Span.
- CAESALPINACEAE
Bauhinia racemosa Lamk.
Cassia absus Linn.
C. auriculata Linn.
C. occidentalis Linn.
Cassia pumila Lamk.
C. siamea Lamk.
C. tora Linn.
Delonix elata (Linn.) Gamble
Tamarindus indica Linn.
- MIMOSACEAE
Acacia jacquemontii Benth.
A. leucophloea (Roxb.) Willd.

Acacia nilotica (Linn.) Del. ssp. *indica* (Benth.) Bremek
A. pennata (Linn.) Willd.

A. senegal Willd.

Albizia lebbeck (Linn.) Benth.

Dichrostachys cinerea (Linn.) W. & Arn.

Mimosa humilis Willd.

Neptunia oleracea Lour.

Pithecellobium dolce (Roth.) Benth.

Prosopis cineraria (Linn.) Drace

P. juliflora (Sw.) DC.

VAHILLACEAE

Vahlia digyna (Retz.) O. Kuntze

COMpositae

Anogeissus sericea Benth.

Combretum ovalifolium Roxb.

Quisqualis indica Linn.

Terminalia bellirica (Gaertn.) Roxb.

T. catappa Linn.

T. crenigluta Roxb.

MYRTACEAE

Eucalyptus sp.

Psidium guajava Linn.

Syzygium hirsutum Wall. ex Gamble

LYTHRACEAE

Anmannia baccifera Linn.

A. multiflora Roxb.

A. senegalensis Lamk.

Lawsonia inermis Linn.

Woodfordia fruticosa (Linn.) Kurz

PLUMACEAE

Funica granatum Linn.

CINNABARACEAE

Ludwigia adscendens (Linn.) Harv.

L. octovalvis ssp. *sessiliflora* (Michx.) Raven

L. perennis Linn.

TRAPACEAE

Trapa natans Linn. var. *bipinnosa* (Roxb.) Makino

CARIACACEAE

Carica papaya Linn.

CUCURBITACEAE

Citrullus colocynthis (Linn.) Schrad.

Coccinia grandis (Linn.) Voigt

Cnidoscolus cerasiformis (Stocks) Hook. f.

Cucurbita callosa (Rottl.) Cogn. ex Cogn. & Harms

Lagenaria leucantha (Duch.) Rusby

Luffa acutangula (Linn.) Roxb.

L. cylindrica (Linn.) M. J. Roem.

Monardica dioica Roxb. ex Willd.

Mukia maderaspatana (Linn.) M. Roem.

Trichosanthes bracteata (Lamk.) Voigt

T. cucumerina Linn.

CUCURBITACEAE

Opuntia elatior Mill.

ALOACEAE

Trianthema portulacastrum Linn.

T. triquetra Roul. ex Willd.

MOLLUGINACEAE

Glinus lotoides Linn.

G. oppositifolius (Linn.) DC.

Mollugo nudicaulis Lamk.

APiales (UMBELLIFERAE)

Anethum graveolens Linn.

Coriandrum sativum Linn.

Foeniculum vulgare Mill.

ALANGIACEAE

Alangium salvifolium (Linn. f.) Wang

RUBIACEAE

Borreria articulata (Linn. f.) F. N. Will

B. stricta (Linn. f.) Schult.

Dentella repens (Linn.) Forst.

Gardenia resinifera Roth

Ixora brachiata Roxb.

Mirugyna parviflora Kurth.

Morinda tomentosa Heyne ex Roth

Oldenlandia corymbosa Linn.

Xerophyllum spinosum (Thunb.) Ker

X. oliginosa (Retz.) Malleshw.

ASTERACEAE (COMPOSITAE)

Acanthospermum hispidum DC.

Amberboa ramosa (Roxb.) Jafri

Bidens biternata (Linn.) Merr. & Sheriff

Blainvillea acmella (Linn.) Phil.

Blumea bifoliata DC.

B. eriantha DC.

B. membranacea DC.

B. obliqua (Linn.) Drace

B. oxyodonta DC.

Caesulia axillaris Roxb.

Dicoma tomentosa Cass.

Echinops echinatus Roxb.

Eclipta prostrata (Linn.) Linn.

Emilia sonchifolia (Linn.) DC.

Glossocardia boswelliae (Linn. f.) DC.

Gnaphalium indicum Linn.

G. pulvinatum Del.

Grangea madagascariensis Poir.

Legnera aurita (Linn. f.) Sch-Bip.

Launaea fallax (Jacq. & Spach.) O. Kuntze

- Sonchus asper* (Linn.) St. Hil.
Sphaeranthus indicus Linn.
Tridax procumbens Linn.
Vernonia cinerea (Linn.) Less.
Vicoa indica (Willd.) DC.
Xanthium strumarium Linn.

LORELIAEAE

- Lobelia alsinoides* Lamk.

PRIMULACEAE

- Antigallis arvensis* Linn. var. *cineraria* Gren. & Godr.

RAPONTEAE

- Madhuca indica* J. F. Gmel.
Manilkara hexandra (Roxb.) Hubb.

EBENACEAE

- Diospyros melanoxylon* Roxb.

SALVADORACEAE

- Salvadora neoides* Decne.

- S. persica* Linn.

APOCYNACEAE

- Carissa congesta* Wt. & Arn.

- Catharanthus pusillus* G. Don

- C. roseus* (Linn.) G. Don

- Holarhena antidysenterica* (Linn.) Wall. ex A. DC.

- Nerium indicum* Mill.

- Thevetia peruviana* (Pers.) Merr.

- Wrightia tinctoria* R. Br.

ASCLEPIADACEAE

- Calotropis procera* (Ait.) R. Br.

- Ceropegia bulbosa* Roxb.

- Drogea volubilis* (Linn.) Benth. ex Hook. f.

- Leptadenia reticulata* (Retz.) Wt. & Arn.

- Pentatropis spiralis* (Forsk.) Decne.

- Pergularia daemia* (Forsk.) Choisy

- Telosma pallida* (Roxb.) Craib.

PERIPLOCACEAE

- Hemidiodia indicus* (Linn.) Schulz.

GENTIANACEAE

- Canscora diffusa* (Vahl) R. Br.

- Endostema hyssopifolium* (Willd.) Verdc.

- Exacum pedunculatum* Linn.

- Hoppea dichotoma* Willd.

- Nymphoides cristatum* (Roxb.) O. Kuntze

- N. indicum* (Linn.) O. Kuntze

HYDROPHYLACEAE

- Hydroclea zeylanica* (Linn.) Vahl

BORAGINACEAE

- Heliotropium marifolium* Retz.

- H. marifolium* Retz. var. *laxiflora* Cooke

- H. ovalifolium* Forsk.
H. subulatum Hochst. ex DC.
H. supinum Linn.
Trichodesma amplexicaule Roth
T. zeylanicum (Burm. f.) R. Br.

ERETIACEAE

- Coldenia procumbens* Linn.
Ehretia aspera Roxb.
E. laevis Roxb.

CONVOLVULACEAE

- Convolvulus microphyllus* (Roth) Sieb.

- Evolvulus alsinoides* (Linn.) Linn.

- Hewittia sublobata* (Linn. f.) O. Kuntze

- Ipomoea aquatica* Forsk.

- I. batatas* (Linn.) Lamk.

- I. dichroa* (Roem. & Schult.) Choisy

- I. eriocarpa* R. Br.

- I. moricata* (Linn.) Jacq.

- I. nil* (Linn.) Roth

- I. obscura* (Linn.) Ker-Gawl.

- I. pes-tigridis* Linn.

- I. quamoleti* Linn.

- I. sepiaria* Koen. ex Roxb.

- I. sindica* Stapf

- I. sinensis* (Desr.) Choisy

- I. triloba* Linn.

- Merremia neglecta* (Linn.) Urb.

- M. gangetica* (Linn.) Cufod.

- M. tridentata* (Linn.) Hall. F.

- Operculina turpethum* (Linn.) Silva-Manso

- Rivea hypocrateiformis* Choisy

CUSCUTACEAE

- Cuscuta chinensis* Lamk.

- C. reflexa* Roxb.

SOLANACEAE

- Capsicum annum* Linn. var. *acuminatum* Fingerh.

- Datura metel* Linn.

- Lycopersicon lycopersicum* (Linn.) A. Siliaw.

- Nicotiana tabacum* Linn.

- Physalis minima* Linn.

- Solanum melongena* Linn.

- S. nigrum* Linn.

- S. surattense* Burm. f.

- Withania somnifera* (Linn.) Dun.

SCROPHULARIACEAE

- Bacopa monnierii* (Linn.) Pers.

- Dopatrium junceum* (Roxb.) Buch-Ham. ex Benth.

- Glossostigma diandrum* (Linn.) O. Kuntze

- Limnophila indica* (Linn.) Druce

- Linnophila sessiliflora* (Vahl) Blume
Lindenbergia muraria (Roxb.) P. Bischl
Lindernia ciliata (Colm.) Penn.
L. crustacea (Linn.) F. Muell.
L. oppositifolia (Retz.) Muk.
L. parviflora (Roxb.) Haines
Sepedea delphinifolia (Roxb.) G. Don
Stemodia viscosa Roxb.
Sruga angustifolia (Don) Sald.
S. generoides (Willd.) Varke var. *minor* Sant.
Verbascom chinense (Linn.) Sant.
- OBORANGIACEAE
- Orobanchus cernua* Loef. var. *nepalensis* DC.
- LENTIBULARIACEAE
- Utricularia inflata* Forsk. var. *stellaris* Taylor
- BIGNONIACEAE
- Dolichandra falcata* Seem. var. *lawii* (Seem.)
Haines
- Millingtonia hortensis* Linn. f.
- PEDALIACEAE
- Scammonia indicum* Linn.
- MARTYNIACEAE
- Martynia annua* Linn.
- AGANTHACEAE
- Ahatoda vaseca* Nees
Bartsia prattensis Sant.
B. prionitis Linn.
Bipharis maderaspaspatina (Linn.) Roth
B. repens (Vahl) Roth
Dicliptera verticillata (Forsk.) Christens.
Dipteronanthus prostratus (Poir.) Nees
Elytraria acuminis (Linn. f.) Lindau
Hemidiodiaulus polyspermus Nees
Hygrophila auriculata (Schlecht.) Heine
Lepidagathis trinervis Wall. ex Nees
Nearanthus sphaerostachyus (Nees) Dalz.
Peristrophe bicalycata (Retz.) Nees
Rungia repens (Linn.) Nees
- VERBENACEAE
- Clerodendrum multiflorum* (Burm. f.) O. Kuntze
Gmelina arborea Roxb.
Lastrea canara Linn. var. *aculeata* (Linn.) Mold.
Phyla nodiflora (Linn.) Greene
Tectea grandis Linn. f.
Vitea negundo Linn.
- LAMIACEAE (LABIATAE)
- Anisomeles indica* (Linn.) O. Kuntze
Hypia suaveolens (Linn.) Poit.

- Leucas aspera* (Willd.) Spreng.
L. biflora R. Br.
L. longiflora Benth.
Moschomia polystachyum (Linn.) Benth.
Ocimum basilicum Linn.
O. canum Linn.
O. sanctum Linn.

NYCTAGINACEAE

- Boehravia chinensis* (Linn.) Druec.
B. diffusa Linn.
Bougainvillea glabra Choisy

AMARANTHACEAE

- Achyranthes aspera* Linn.
Aerva lanata (Linn.) Juss.
Alternanthera sessilis (Linn.) DC.
Amaranthus hybridus Linn. spp. *cruentus* (Linn.)
Thell. var. *paniculatus* (Linn.) Thell.
A. spinosus Linn.
A. tricolor Linn.
A. viridis Linn.
Celosia argentea Linn.
Digera muricata (Linn.) Mart.
Gomphrena celosioides Mart.
Nothosperma brachista (Linn.) W.C.
Pupalia lappacea (Linn.) Juss.

CHENOPodiaceae

- Chenopodium album* Linn.
C. murale Linn.

POLYGONACEAE

- Polygonum plebeium* R. Br.
 LORANTHACEAE
Dendrophthoe falcata (Linn. f.) Etting.

EUPHORBIACEAE

- Acalypha indica* Linn.
Bridelia squamosa (Muell.-Arg.) Gehr.
Chuorophora prostrata Dalz.
Euphorbia hirta Linn.
E. nerifolia Linn.
E. orbiculata H. B. K.
E. parviflora Linn.
E. prostrata Ait.
E. thymifolia Linn.
E. tirucalli Linn.
Jatropha curcas Linn.
J. gossypifolia Linn.
Kirganelia reticulata (Poir.) Baill.
Phyllanthus fraternus Webster
P. simplex Retz.

- 524
- Ricinus communis* Linn.
Securigera varia (Roth ex Willd.) Pax & Hoffm.
- ULMACEAE
- Holopeltis integrifolia* (Roth) Planck
- MORACEAE
- Ficus amplissima* Sonn.
F. benghalensis Linn.
F. hispida Linn. f.
F. racemosa Linn.
F. religiosa Linn.
Streblus asper Linn.
- CERATOPHYLLACEAE
- Ceratophyllum demersum* Linn.
- HYDROCHARITACEAE
- Hydrilla verticillata* (Linn. f.) Royle
- Vallisneria spiralis* Linn.
- HYPOXIDACEAE
- Curculigo orchioides* Gaertn.
- DISCOREACEAE
- Dioscorea bulbifera* Linn.
- LILIACEAE
- Allium cepa* Linn.
A. sativum Linn.
Aloe barbadensis Mill.
Asparagus officinalis Bak.
A. racemosus Willd. var. *javanicus* (Kunth) Bak.
Asphodelus tenuifolius Cav.
Gloriosa superba Linn.
- PONTEDERIACEAE
- Monochoria vaginalis* (Burm. f.) Presl
- COMMELINACEAE
- Commelinia benghalensis* Linn.
C. diffusa Burm. f.
C. forskalaei Vahl
C. hastarilis Clarke
C. kurzii Clarke
Cyanotis axillaris (Linn.) Schult.
Murdannia nudiflora (Linn.) Breitw.
M. semiteres (Dals.) Sant.
- ARECACEAE (PALMAE)
- Borassus flabellifer* Linn.
Phoenix sylvestris Roxb.
- TYPHACEAE
- Typha angustata* Bory & Chah.
 LEMNACEAE
- Lemna paucicostata* Hegelm.
Wolffia arrhiza (Linn.) Wimm.
W. microscopica (Griff.) Kurz
- ALISMACEAE
- Limnophyton obtusifolium* (Linn.) Miq.
- NAJADACEAE
- Najas minor* All.
N. kurziana Rendle
- APONOGETONACEAE
- Aponogeton natans* (Linn.) Engl. & Krause
- BUTOMACEAE
- Tenagogcharis latifolia* (D. Don) Buchen.
- ERIOGAULACEAE
- Eriocaulon elatior* Fyson
- CYPERACEAE
- Bulbostylis barbata* (Roth.) Kunth
Cyperus alopecuroides Rottb.
C. bulbosus Vahl
C. compressus Linn.
C. difformis Linn.
C. iria Linn.
C. lievigatus Linn.
C. michelianus (Linn.) Link. ssp. *pygmaeus* (Roth.) Achers. & Griseb.
C. nutans Vahl var. *clematoides* (Kunth) Haines
C. pangorei Rottb.
C. pseudokyllingioides Kükenth.
C. pumilus Linn.
C. pumilus Linn. var. *membranaceus* (Vahl) Clarke
C. rotundus Linn.
C. squarrosum Linn.
C. tenuispica Steud.
C. triceps (Roth.) Endl.
Eleocharis atropurpurea Kunth
E. dulcis (Burm. f.) Triana ex Henschel.
E. geniculata (Linn.) Roem. & Schult.
Fimbristylis argentea (Roth.) Vahl
F. dichotoma (Linn.) Vahl
F. falcatu (Vahl) Kunth
F. ferruginea (Linn.) Vahl
F. littoralis Gaud.
F. miliacea (Linn.) Vahl
F. ovata (Burm. f.) Kern
F. schoenoides (Retz.) Vahl
F. sieberiana Kunth
F. tenera Roem. & Schult.
Fuirena ciliaris (Linn.) Roxb.
F. tuvensis Deshpande & Shah
Scirpus articulatus Linn.
S. brachyceras Hochst. ex A. Rich
S. kyllingioides Boeck.
S. lateriflora Gmel.
S. littoralis Schrad. var. *subulatus* (Vahl) Chiov.

Sirpus maritimus Linn.
S. rojoi (Nees) Parker
S. squarrosum Linn.

POACEAE (GRAMINEAE)

- Alstroemeria cimicina* (Linn.) Stapf
Apluda mutica Linn.
Aristida adscensionis Linn. var. *adscensionis*
Bracharia ramosa (Linn.) Stapf
Capillipedium hugelii (Hark.) Stapf
Cochlosperma biforme Roxb.
Cleistis barbata (Linn.) Sw.
C. dolichostachya Lagas.
Cymbopogon martinii (Roxb.) Wats.
Cynodon dactylon (Linn.) Pers.
Dactyloctenium aegyptium (Linn.) P. Beauv.
D. indicum Boiss.
Desmostachya bipinnata (Linn.) Stapf
Dianthus annulatum (Forsk.) Stapf
Dineira retroflexa (Vahl) Panz.
Echinocloa column (Linn.) Link.
E. frumentacea Link.
Engrostis ciliensis (Linn.) Vig.-Lut.
E. ciliaris (Linn.) R. Br. var. *brachystachya* Boiss.
E. ciliaris (Linn.) R. Br. var. *charkei* Stapf ex Hook. f.
E. diarrhoea (Schult.) Steud.
E. japonica (Thunb.) Trin.
E. pilosa (Linn.) P. Beauv.
E. tremula (Linn.) P. Beauv. ex Roem. & Schult. var.
tenuis
E. tremula Hochst. ex Steud.
Eremopogon foeculatus (Del.) Stapf
Festuca trispicata (Schult.) Hene.
Hakelochloa granularis (Linn.) O. Kuntze
Heteropogon contortus (Linn.) Beauv. ex Roem. &
Schult.
Ichneumon rugosum Salisb.
Melanocenchrus jacquemontii Juss. & Spach.
Oryza nivara Sharma & Shastri
O. sativa Linn.
Panicum antidotale Retz.
P. milletorum Linn.
P. pilosum Trin.

- Paspalidium flavidum* (Retz.) A. Camus
P. geminatum (Forsk.) Stapf
Paspalum scrobiculatum Linn.
Pennisetum typhoides (Burm. f.) Stapf & Hobb.
Pteris indica (Linn.) O. Kuntze
Phragmites kurka (Retz.) Trin. ex Steud.
Pseudoraphis spinosces (R. Br.) Vickery
Saccharum officinarum Linn.
S. spontaneum Linn.
Setaria glauca (Linn.) P. Beauv.
S. tomentosa (Roxb.) Kunth
S. verticillata (Linn.) P. Beauv.
Sorghum vulgare Pers.
Sporobolus coromandelianus (Retz.) Kunth
S. diander (Retz.) P. Beauv.
Tetrapogon tenellus (Roxb.) Chiov.
Themeda quadrivalvis (Linn.) O. Kuntze var.
quadrivalvis
Tragus biflorus (Roxb.) Schult.
Triticum aestivum Linn.
Vetiveria zizanioides (Linn.) Nash
Zea mays Linn.

REFERENCES

- BEST, S. J. Flora study of Ratnagiri and surrounding hills. Ph.D. Thesis, 1958. Not seen.
- , S. D. SAHOO AND R. P. BHATE. Additions to the flora of Pavagadh hill, Gujarat State. J. Bot. Soc. Ind. 86, No. 5(2): 512-524, 1965.
- CHAVAN, S. R. AND G. M. DIA. The flora of Pavagadh hill, M. S. Univ. Baroda. Bot. Mem. No. 1: 1-296, 1965.
- , AND S. J. BEST. New plant records from cravahile Bombay State III. J. Bot. Soc. Ind. Mem. No. 5(2): 779-781, 1965.
- CHETTRI, J. G., G. L. SHAH AND M. B. DESHPANDE. *Elaeis obovata* Willd.: A new record for central India. Bull. Bot. 64: 143-146, 1967.
- , AND G. L. SHAH. Some more plants from Pavagadh. Ibid. 66(2): 463-466, 1969.
- DESHPANDE, M. B. AND G. L. SHAH. A new species of *Palaeo* from Gujarat, India. Bull. Bot. Surv. India 10(2): 289-293, 1968.
- SAXTON, W. T. Additional notes on plants of Northern Gujarat. J. Bot. Soc. Ind. 9(2): 273-282, 1922.
- SHAH, G. L. AND M. B. DESHPANDE. Additions to the flora of Bombay. J. Bot. Soc. Ind. Mem. No. 5(1): 209-222, 1965.
- , AND J. A. DHAMKAR. Further contribution to the flora of Pavagadh hill near Baroda, Gujarat. Bull. 67(2): 279-294, 1965.